



Evidence-Based Evaluation of Complementary Health Approaches for Pain Management in the United States

Richard L. Nahin, PhD, MPH; Robin Boineau, MD, MA; Partap S. Khalsa, DC, PhD; Barbara J. Stussman, BA; and Wendy J. Weber, ND, PhD, MPH



From the National Center for Complementary and Integrative Health, National Institutes of Health, Bethesda, MD.

CME Activity

Target Audience: The target audience for *Mayo Clinic Proceedings* is primarily internal medicine physicians and other clinicians who wish to advance their current knowledge of clinical medicine and who wish to stay abreast of advances in medical research.

Statement of Need: General internists and primary care physicians must maintain an extensive knowledge base on a wide variety of topics covering all body systems as well as common and uncommon disorders. *Mayo Clinic Proceedings* aims to leverage the expertise of its authors to help physicians understand best practices in diagnosis and management of conditions encountered in the clinical setting.

Accreditation: Mayo Clinic College of Medicine is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Statement: Mayo Clinic College of Medicine designates this journal-based CME activity for a maximum of 1.0 AMA PRA Category 1 Credit(s).™ Physicians should claim only the credit commensurate with the extent of their participation in the activity.

MOC Credit Statement: Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

Learning Objectives: On completion of this article, you should be able to (1) name complementary health approaches used for pain management; (2) discuss the evidence supporting the use of complementary health approaches for pain management; and (3) provide examples to your patients of complementary health approaches that might be considered as part of a comprehensive pain management plan.

Disclosures: As a provider accredited by ACCME, Mayo Clinic College of Medicine (Mayo School of Continuous Professional Development) must

ensure balance, independence, objectivity, and scientific rigor in its educational activities. Course Director(s), Planning Committee members, Faculty, and all others who are in a position to control the content of this educational activity are required to disclose all relevant financial relationships with any commercial interest related to the subject matter of the educational activity. Safeguards against commercial bias have been put in place. Faculty also will disclose any off-label and/or investigational use of pharmaceuticals or instruments discussed in their presentation.

Disclosure of this information will be published in course materials so that those participants in the activity may formulate their own judgments regarding the presentation.

In their editorial and administrative roles, William L. Lanier, Jr, MD, Terry L. Jopke, Kimberly D. Sankey, and Nicki M. Smith, MPA, have control of the content of this program but have no relevant financial relationship(s) with industry. The authors report no competing interests.

Method of Participation: In order to claim credit, participants must complete the following:

1. Read the activity.
2. Complete the online CME Test and Evaluation. Participants must achieve a score of 80% on the CME Test. One retake is allowed.

Visit www.mayoclinicproceedings.org, select CME, and then select CME articles to locate this article online to access the online process. On successful completion of the online test and evaluation, you can instantly download and print your certificate of credit.

Estimated Time: The estimated time to complete each article is approximately 1 hour.

Hardware/Software: PC or MAC with Internet access.

Date of Release: 9/1/2016

Expiration Date: 8/31/2018 (Credit can no longer be offered after it has passed the expiration date.)

Privacy Policy: <http://www.mayoclinic.org/global/privacy.html>

Questions? Contact dletcsupport@mayo.edu.

Abstract

Although most pain is acute and resolves within a few days or weeks, millions of Americans have persistent or recurring pain that may become chronic and debilitating. Medications may provide only partial relief from this chronic pain and can be associated with unwanted effects. As a result, many individuals turn to complementary health approaches as part of their pain management strategy. This article examines the clinical trial evidence for the efficacy and safety of several specific approaches—acupuncture, manipulation, massage therapy, relaxation techniques including meditation, selected natural product supplements (chondroitin, glucosamine, methylsulfonylmethane, S-adenosylmethionine), tai chi, and yoga—as used to manage chronic pain and related disability associated with back pain, fibromyalgia, osteoarthritis, neck pain, and severe headaches or migraines.

© 2016 Mayo Foundation for Medical Education and Research ■ *Mayo Clin Proc.* 2016;91(9):1292-1306

The most recent national estimate suggests that 126 million adults experience some pain in a given year,¹ with about one-third (40 million adults) having severe pain. Pain is often associated with poor general health, health-related disability, and increased health care utilization.² Yet according to the Institute of Medicine,² pain is routinely undertreated in health care; pain care that is provided is often fragmented, without a comprehensive assessment or treatment plan, and patients may encounter difficulty obtaining the full range of potential treatments.

Back pain, joint pain, neck pain, and headaches are among the most common types of pain experienced by US adults (Table 1).³⁻⁶ The prevalence rates for these conditions have remained relatively stable over time. Among the many pharmacological and nonpharmacological approaches that have been incorporated into pain management strategies are complementary health approaches. This broad category of care includes procedures by licensed practitioners such as acupuncturists, chiropractors, and massage therapists, as well as self-care approaches such as relaxation techniques (eg, meditation) and meditative movement-based approaches (eg, yoga and tai chi) and natural products such as glucosamine and herbal medicines. National surveys going back more than 25 years have consistently found that these complementary approaches are used by about 30% to 40% of the US public in a given year,⁷⁻¹¹ although use of a given approach may wax and wane over time (Table 2).

Although a substantial part of this use is for overall wellness and prevention,^{12,13} painful conditions are the most common health

TABLE 1. Age-Adjusted Percentages of Selected Health Conditions Among US Adults, NHIS 2002, 2007, 2012^a

Health condition	2002 ³	2007 ⁴	2012 ^{5,6}
Low back pain in the past 3 mo	26.4%	25.4%	27.6%
Arthritis	20.9%	20.3%	20.6%
Neck pain in the past 3 mo	13.8%	12.8%	13.9%
Severe headache or migraine in the past 3 mo	15.0%	12.3%	14.1%
Fibromyalgia during lifetime	NA	NA	1.75%

^aNA = not available; NHIS = National Health Interview Survey.

TABLE 2. Age-Adjusted Percentages of Use for Selected Complementary Health Approaches by US Adults, 2002-2012

Complementary health approach	2002 ⁷	2007 ⁸	2012 ⁹
Acupuncture	1.1%	1.4%	1.5%
Manipulation	7.5%	8.6%	8.4%
Massage therapy	5.0%	8.3%	6.9%
Meditation	7.6%	9.4%	8.0%
Natural product supplements	18.9%	17.7%	17.7%
Yoga, tai chi, and qigong	5.8%	6.7%	10.1%

conditions for which individuals turn to these complementary approaches.^{7,8,10,11,14} In 2007, for example, about 14.3 million adults used a complementary health approach for their back pain, about 5.0 million used these approaches for their neck pain, and 3.1 million for their arthritis.⁷ Far fewer individuals used complementary health approaches for other chronic diseases such as depression (1.0 million), hypertension (0.8 million), diabetes (0.7 million), or cancer (0.4 million).

Based on national survey data,¹⁴ this high use of complementary health approaches for painful conditions translated into \$8.5 billion in out-of-pocket payments for these approaches to manage back pain, \$3.6 billion to manage neck pain, and \$2.3 billion to manage arthritis. Substantially less is spent out-of-pocket on complementary health approaches to treat other chronic health conditions such as depression (\$1.1 billion), hypertension (\$0.7 billion), diabetes (\$0.3 billion), and cancer (\$0.2 billion).

Given the high use of complementary health approaches for pain, a number of specific complementary approaches have undergone mechanistic and clinical evaluations culminating in phase 3 trials. This article examines the clinical trial evidence for the efficacy and safety of several widely used approaches—acupuncture, manipulation, massage therapy, relaxation techniques including meditation, selected natural product supplements (eg, chondroitin and glucosamine), tai chi, and yoga (defined in Supplemental Appendix 1, available online at <http://www.mayoclinicproceedings.org>)—as used to manage chronic pain and related disability associated with back pain, osteoarthritis (OA), neck pain, and severe headaches or migraines, conditions frequently seen and managed in the primary care setting.

Fibromyalgia was included in this review as an example of a complex pain syndrome that is often managed with a multimodal approach that may include complementary approaches. Cancer pain is certainly a major public health concern but is more likely to be addressed outside the primary care setting (eg, by oncologists, at cancer centers, as part of palliative care).

The randomized, controlled clinical trial (RCT) is considered the strongest study design for investigating the efficacy and safety of pharmacological, behavioral, and physical interventions. To identify examples of RCTs for each complementary approach, we performed searches of the MEDLINE database for articles published from January 1, 1966, through March 31, 2016, using the search strategy outlined in [Supplemental Appendix 2](#) (available online at <http://www.mayoclinicproceedings.org>). In order to make this review as relevant as possible to primary care physicians in the United States, we limited this review to RCTs either performed in the United States or that included participants from the United States. This decision was based on 2 factors. First, the unique health care system in the United States vs other countries means that the standard care or usual care control groups used in the United States and other countries may vary substantially. Thus, whether a given complementary approach performs better than usual care in another country may not reflect how the approach would perform in US trials. Another factor is that the training and licensure of acupuncturists, chiropractors, and naturopathic doctors vary substantially among countries, as does the marketing, regulation, and use of dietary supplements. For instance, in Germany, the location of some of the largest acupuncture trials, acupuncture is only practiced by medical doctors, whereas the vast majority of acupuncture treatment in the United States is provided by licensed acupuncturists. Thus, the findings from German trials may not be directly comparable to acupuncture as practiced in the United States.

Brief summaries of the reviewed RCTs are presented with details provided in online supplemental tables. The findings of these RCTs also illustrate several methodological issues that should be considered when interpreting the trial data. These issues are summarized briefly at the end of this article.

LOW BACK PAIN

Acupuncture

We found 4 RCTs (total participants, 1092)¹⁵⁻¹⁸ that assessed the clinical benefit of acupuncture for treatment of low back pain (LBP) (age range, 28-60 years; most participants were white) and used primary study outcomes of self-report of pain intensity (numeric rating scale or visual analog scale [VAS]) and/or functional disability (Roland-Morris Disability Questionnaire, Oswestry Disability Index [ODI], or Disability Rating Index). Cherkin et al^{15,16} reported modest improvement in pain intensity and function compared with usual care. In pregnant women using auricular acupuncture, Wang et al¹⁷ found a significant reduction in pain intensity and improved functional status compared with no treatment. Comparison of verum to sham acupuncture had mixed results, with 2 RCTs^{16,18} finding no significant difference and 1 RCT¹⁷ finding a slight but significant difference. No significant adverse events were reported.

Massage Therapy

We identified 8 RCTs that studied the use of massage for LBP^{15,19-25} (total participants, 829). Massage types included Swedish/relaxation, structural, structural integration, and muscle energy; sessions varied in duration from 15 to 90 minutes. For chronic LBP (cLBP), 2 larger studies^{15,19} comparing massage with usual care reported modest improvements in pain and function at 10 weeks, but the benefit was not sustained at 52 weeks.¹⁵ Three smaller studies compared massage to either usual outpatient rehabilitation²⁴ or relaxation^{22,23} and did not observe significant between-group differences for pain and/or function. For acute or subacute LBP, 2 smaller studies found significant, albeit modest, improvements in pain compared with no treatment (−1.5 points on a numeric rating scale)²¹ or function (−18% on the ODI)²⁵ compared with a putative placebo. No RCTs reported significant adverse events.

Osteopathic Manipulative Therapy

Six RCTs of osteopathic manipulative therapy (OMT) for LBP were identified (total participants, 1308).²⁶⁻³¹ Two RCTs examined OMT compared with sham OMT for cLBP^{30,31} using

similar intervention paradigms and reported mixed results, with 1 finding no significant difference and 1 reporting a 9-mm reduction in pain intensity on a VAS. Two RCTs compared OMT with usual care for acute or subacute LBP,^{26,27} and both reported no significant improvement for function/disability but mixed results for pain intensity for between-group differences. In pregnant women with LBP, 2 studies compared the benefit of adding OMT or placebo ultrasound treatment to usual obstetric care and reported significant modest improvements with added care but no significant between-group differences.^{28,29} No RCTs reported significant adverse events.

Spinal Manipulation

We reviewed 24 RCTs³²⁻⁵⁵ (total participants, 4503; all adults) of spinal manipulation (SM) for LBP. Recent data suggest that for cLBP, the “dose” of SM (defined as the number of sessions) may affect outcomes,^{45,46,56} and hence the dose utilized was dichotomized for this report depending on whether 6 or more sessions of SM were provided in a given study. No RCTs reported significant adverse events.

Chronic LBP. There were 9 studies of SM for cLBP (total participants, 1882),^{32,35-37,40,44-46,50} 8 of which employed 6 or more sessions for treatment, and 1 of these studies included principally (more than two-thirds) adult women.³² Among the 4 larger RCTs (each with 200 or more participants)^{37,44,46,50} that compared high-velocity low-amplitude SM with an active comparison (exercise, usual care, physical therapy, light touch), 3 found significant, albeit modest, between-group differences for pain intensity and/or function. Two studies directly examined the dose of high-velocity low-amplitude SM,^{45,46} and both found that higher doses (12 and 18 sessions of SM) provided larger improvement at 12 and 52 weeks, respectively. One RCT utilized a putative placebo SM³² and found that after the first session, the results of verum SM were better than sham SM and better than no treatment, but at 2 weeks, there were no significant between-group differences. One RCT compared thrust to non-thrust SM⁴⁰ (less than 6 sessions) and reported no significant between-group differences.

Acute, Subacute, or Mixed LBP. There were 15 studies of SM for either acute, subacute,

or a mixture of types of LBP (total participants, 2621).^{33,34,38,39,41-43,47-49,51-55} Six of these studies were of modest to moderate size (ie, >100 participants) and utilized 6 or more sessions of SM.^{38,42,49,51,53,54} The results of these studies were mixed, with some reporting modest significant benefit of SM compared with active intervention (physical therapy, education [“back school”], medication, usual care) at about 4 weeks for pain intensity and/or function,^{42,49,54} but others reporting no significant between-group differences.^{38,51,53} One RCT examined pregnant women with LBP⁴² and found that adding SM and exercise to usual obstetric care provided modest improvement in pain and function/disability.

There were 5 studies of SM for either acute and/or subacute LBP that used less than 6 treatment sessions,^{33,39,41,47,55} but only 1 of these (and the largest) was an effectiveness study: Fritz et al⁴¹ conducted an RCT of 4 sessions over 4 weeks of SM plus exercise vs usual primary care and at 3 months found significant between-group improvement in function (−3.2 points on the ODI). The other 4 studies (sample sizes ranged from 54 to 123 patients) compared different types of SM and/or SM with an active intervention (eg, exercise). In general, these 4 studies reported that all groups had improvement in back pain, and there were very small or no significant between-group differences. However, it is unclear whether these 4 studies were sufficiently powered to definitively ascertain whether differences existed.

There were 4 studies of SM that by design addressed mixed LBP^{34,48,52,53} (ie, the inclusion criteria allowed participants with acute, subacute, or chronic LBP); 3 of these studies utilized 6 or more treatment sessions. All studies had one or more active comparison groups (eg, massage, electrical stimulation, usual care, corticosteroid injection, physical therapy), and in all studies, all groups improved but there were no significant between-group differences.

Yoga

We identified 6 RCTs of yoga for cLBP⁵⁷⁻⁶² (total participants, 596; all adults, predominantly female). Three named forms of yoga were studied: hatha,^{57,58} viniyoga,^{59,60} and iyengar^{61,62}, all were performed in group settings, with class durations from 60 to 90 minutes and the

number of sessions ranging from 12 to 24, either once or twice per week, with recommendations for home practice. Compared with usual care, 2 studies^{59,60} found that yoga provided improvements in pain and function, but the results were mixed when compared with exercise/stretching. A dose-response study⁵⁷ compared once-weekly to twice-weekly classes and found that they produced equivalent improvements in pain intensity and function. Three smaller studies compared yoga with wait list^{58,61} or education control⁶² and reported significant modest reductions in pain intensity and function/disability. No RCTs reported significant adverse events.

Additional information on all back pain RCTs can be found in [Supplemental Table 1](#) (available online at <http://www.mayoclinicproceedings.org>).

FIBROMYALGIA

All trials we reviewed for fibromyalgia used the 1990 American College of Rheumatology (ACR) classification criteria,⁶³ except one⁶⁴ that used an older definition.

Acupuncture

Four RCTs examined acupuncture vs sham acupuncture for pain, physical function, global well-being, sleep, fatigue, and adverse events.⁶⁵⁻⁶⁸ Martin et al⁶⁸ found a significant improvement between electroacupuncture vs sham electroacupuncture. Differences were seen on the Fibromyalgia Impact Questionnaire (FIQ) scores for fatigue and anxiety. No other trial found significant differences between groups on any outcome. There were no serious adverse events reported in any of these studies. In one study, minor adverse events (eg, discomfort at the site of needle insertion or simulation of needles) were reported by 89% of participants.⁶⁵

Relaxation Techniques

Two studies (93 total participants, mostly female and white) investigated biofeedback vs control groups (attention control and placebo) as a treatment for symptoms of fibromyalgia.^{64,69} Buckelew et al⁶⁴ found a significant improvement in the Tender Point Index score in the biofeedback group vs an attention control group but not for any other outcome measure. Nelson et al⁶⁹ did not find any differences between

biofeedback and a placebo biofeedback. A small study (90 women) found that mindfulness-based stress reduction significantly reduced perceived stress and sleep disturbance and lessened the severity of symptoms in persons with fibromyalgia vs a wait list control group.⁷⁰ Another RCT examined the effects of affective self-awareness, a technique that places primary importance on the awareness and expression of emotions underlying fibromyalgia symptoms, in 45 women with fibromyalgia and found significant pain reduction and improved physical functioning vs a wait list control group.⁷¹ Astin et al⁷² examined the effects of an intervention combining mindfulness meditation and qigong and found that the combined intervention yielded no better results than an educational/support control group for pain, depression, and physical functioning. Two studies^{73,74} (112 total participants, mostly female) examined guided imagery vs usual care as a treatment for symptoms of fibromyalgia. One study⁷⁴ found a significant decrease in the FIQ score compared with the usual care control group. The second study⁷³ found positive effects of guided imagery on pain intensity, fatigue, and depression vs the control group. Both studies found improvements in self-efficacy for managing symptoms. Only one study⁶⁹ reported on adverse events, and none were noted.

Massage

A small study (12 women) examined Swedish massage vs myofascial release therapy for fibromyalgia symptoms.⁷⁵ No difference was seen between groups on the FIQ.

Tai Chi

A study of 98 adults with fibromyalgia (aged 40 years and older, mostly white and female) compared Yang-style tai chi (modified for fibromyalgia patients) with an educational control and found that the tai chi group had a greater decrease in the FIQ score.⁷⁶ Another study (59 adults with fibromyalgia) compared Yang-style tai chi with a control combining wellness education and stretching classes and found that the tai chi group had greater improvement in the FIQ score.⁷⁷

Yoga

A small study (53 women) investigated yoga vs wait list for management of fibromyalgia

symptoms and found that those practicing yoga had significant improvement in the FIQ score.⁷⁸ No adverse events were noted.

Additional information on all fibromyalgia RCTs can be found in [Supplemental Table 2](#) (available online at <http://www.mayoclinicproceedings.org>).

NECK PAIN

Massage

Four randomized controlled trials examined whether massage could relieve symptoms associated with chronic neck pain.⁷⁹⁻⁸² One study did not report patient demographic characteristics, and the others studied patients aged 20 to 64 years. Primary outcomes included scores on the Neck Disability Index (NDI) (a 10-item neck pain questionnaire), the pain VAS, and range of motion. Sherman et al⁸¹ found significant improvement of the NDI score for those randomized to 10 massage therapy session over 10 weeks vs those assigned to a self-care book on managing neck problems. In an RCT by Field et al,⁸⁰ individuals were randomized to either a wait list control or 30 minutes of massage therapy weekly for 4 weeks combined with daily self-massage. At the completion of the intervention, those assigned to massage therapy had improvements in pain and range of motion compared with the control group. Sherman et al⁸² reported a dose-response relationship between the number and duration of massage sessions per week and improvement in the NDI score and neck pain intensity. The findings indicated that 60 minutes of massage 2 to 3 times per week was significantly better than either 30 or 60 minutes of massage once per week after the 4 weeks of treatment. In a follow-up to the study by Sherman et al,⁸² Cook et al⁷⁹ obtained repeated consent from the participants and randomized them to one additional massage therapy session per week for 6 additional weeks. At the end of treatment, those randomized to the additional sessions had significantly improved pain and function vs those who did not receive the additional sessions; the difference between groups was no longer significant after 14 weeks of follow-up.

Spinal Manipulation

We reviewed 3 randomized trials of SM for neck pain.⁸³⁻⁸⁵ One study assessed manipulation

compared with mobilization with a 2×2×2 factorial design: with or without heat or with or without electrical muscle stimulation⁸⁴; no significant differences in outcomes were seen between groups. Evans et al⁸³ compared SM combined with supervised exercising to supervised exercising alone and also to home exercise. After completion of the 12-week intervention, no difference was seen between SM combined with supervised exercising and supervised exercising alone; however, both these groups had significant improvement in neck pain vs only home exercise. Maiers et al⁸⁵ assessed the efficacy of 3 treatments: (1) SM plus home exercise, (2) supervised rehabilitation exercise plus home exercise, and (3) home exercise alone. Spinal manipulation with home exercise produced significantly better reduction in pain than home exercise alone. No significant difference was seen between SM and home exercise vs supervised rehabilitation exercise plus home exercise.

There was one RCT of manual cervical distraction,⁸⁶ a traction-based therapy with low, medium, and high forces assessed. The goal of the study was to identify a viable sham control. The study end points included pain VAS, NDI score, and a credibility and expectancy questionnaire. The investigators did report benefit in medium- or high-force interventions.

Additional information on all neck pain RCTs can be found in [Supplemental Table 3](#) (available online at <http://www.mayoclinicproceedings.org>).

OA OF THE KNEE

Acupuncture

Four RCTs examined whether acupuncture could relieve symptoms associated with OA of the knee.⁸⁷⁻⁹⁰ These studies used similar definitions of knee OA. Participants were predominantly female, had mean ages between 60 and 65 years, and had knee pain for an average of 9 to 11 years. All studies incorporated either the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) total score or the WOMAC pain subscore as the primary outcomes. In 2 of the trials,^{89,90} no difference was seen between verum acupuncture and sham acupuncture for either the primary or secondary outcome measure. The third trial,⁸⁷ which used a sham control, found significantly better

improvement in both the WOMAC pain subscale and function subscale after 14 weeks of treatment. For the 3 trials that also incorporated either an attention control⁸⁷ or standard care controls,^{88,90} verum acupuncture produced significantly better improvement in primary and secondary outcomes than that observed in the control group. Across trials, adverse events associated with acupuncture were few, with the most common complaints being pain at the needling site and muscle soreness that resolved soon after the treatment session ended.

Glucosamine and Chondroitin

We reviewed 8 RCTs examining the efficacy of 2 dietary supplements, glucosamine and/or chondroitin, in individuals with knee OA.⁹¹⁻⁹⁸ These studies varied substantially in how knee OA was defined, as well as in the primary outcome measures used.

Three different configurations of glucosamine were used: glucosamine hydrochloride (HCl),⁹³⁻⁹⁵ glucosamine sulfate,^{91,92,98} and glucosamine bound to a polymer.⁹⁷ One trial did not identify the configuration of glucosamine.⁹⁶ Doses of glucosamine varied from 1000 mg/d for 6 weeks⁹⁸ to 1500 mg/d for 24 weeks.^{91,95} Chondroitin sulfate was studied in 2 trials at a dose of 1200 mg/d for 6 months^{91,95} and in 1 trial at the same dose but for 8 weeks.⁹³ All studies employed a placebo control group. Participants in 2 trials were exclusively⁹³ or predominantly⁹⁶ males, participants in 4 other trials were predominantly females,^{91,94,95,98} and the remaining 2 trials had a close balance of men and women.^{92,97} The mean age of the trial participants varied from 45 years⁹³ to 72 years.⁹⁵ Mixed results were found in trials comparing glucosamine vs a placebo control with pain relief or functional improvement as the primary outcomes. Three trials found glucosamine superior to placebo,^{92,97,98} and 3 trials found no difference between groups.^{91,94,96}

Mixed results were seen in the 3 trials studying a combination of glucosamine and chondroitin. Leffler et al⁹³ found that a combination of glucosamine HCl and chondroitin sulfate (and manganese ascorbate) was superior to placebo for providing pain relief. In 2 other trials, the combination of glucosamine (either sulfate or HCl) and chondroitin sulfate were no better than placebo for either pain relief or function.^{91,95}

Across trials, adverse events were generally mild (gastrointestinal distress being the most common) with no differences seen between those taking glucosamine or chondroitin and those taking placebo.

Massage Therapy

Two RCTs studied the efficacy of Swedish massage therapy for symptoms associated with OA of the knee.^{99,100} The 2 studies were similar in that both defined knee OA using the ACR criteria,¹⁰¹ required a baseline score of at least 40 on the pain VAS, and included participants who were predominantly white females. In the earlier study,¹⁰⁰ after completing the 8-week intervention, participants in the massage group had significant improvement vs those randomized to usual care in the WOMAC total score as well as in each of the WOMAC subscale scores (pain, function, and joint stiffness). In a dosing study,⁹⁹ individuals were randomized to 1 of 5 groups for 8 weeks: (1) usual care, (2) 240 minutes of massage over 8 weeks, (3) 360 minutes of massage, (4) 480 minutes of massage, and (5) 600 minutes of massage. Only individuals receiving at least 480 minutes of massage therapy (groups 4 and 5) had substantial improvement in the WOMAC total score and the WOMAC pain subscale score vs the usual care group. Across the 2 trials, only one adverse event, discomfort at the knee in one participant, was noted.

Methylsulfonylmethane

One RCT compared methylsulfonylmethane (MSM) (6 g/d for 12 weeks) with a placebo control.¹⁰² Knee OA was based on ACR criteria.¹⁰¹ Outcome measures were the WOMAC subscale scores for pain, function, and stiffness. Individuals randomized to MSM had significant improvement in both the WOMAC pain and functions scale scores. However, the authors cautioned that the differences between groups were small (<20%) and probably not clinically meaningful. No difference in adverse events was seen between groups.

S-Adenoylmethionine

One RCT compared the dietary supplement S-adenoylmethionine (SAME) (1200 mg/d) with celecoxib (200 mg/d for 16 weeks) in a

crossover design that included a 1-week washout period.¹⁰³ Knee OA was defined by ACR criteria,¹⁰² and the study population was primarily female and white. The primary outcomes were pain VAS scores, WOMAC subscale scores for pain, function, and stiffness, and clinician assessments of OA severity. By the end of the trial, no differences were seen between the 2 treatment arms on any of the outcome measures. However, the sample size was insufficient to establish equivalency. Overall, adverse events were less common in those taking SAME than in those taking celecoxib.

Tai Chi

We reviewed 4 RCTs that examined the efficacy of tai chi in patients with confirmed knee OA using various criteria for diagnosis.¹⁰⁴⁻¹⁰⁷ The 4 RCTs had similar study populations, primarily white and female, with mean ages ranging from 65 to 79 years. Mean body mass index (calculated as weight in kilograms divided by height in meters squared) was also similar across the studies, ranging from 27.8 to 30.0 kg/m².

In the study by Hartman et al,¹⁰⁵ participants randomized to a Yang-style tai chi group had significant improvement on the Arthritis Self-Efficacy Scale vs those in the routine care group. The remaining trials compared tai chi to attention controls.^{104,106,107} All found that the tai chi (either Yang style or Sun style) groups did better than the attention control groups on both primary and secondary outcomes. Across trials, the most common adverse event associated with tai chi was minor muscle soreness that resolved after a few days.

Yoga

Park et al¹⁰⁸ completed a quasi-experimental trial with participants randomized to either sitting yoga or Reiki; however, the attention control group was chosen as a convenience sample from other participants meeting the inclusion/exclusion criteria. The primary outcome measures were the WOMAC total score and the WOMAC pain, function, and stiffness subscale scores. The mean age of participants was 80 years, with 68.7% being male. Individuals randomized to yoga had substantially better reduction in the WOMAC

function score than either the Reiki group or the attention control group ($P < .02$). No other differences were seen between the 3 groups. No adverse events were reported for any of the groups in this study.

Additional information on all OA RCTs can be found in [Supplemental Table 4](#) (available online at <http://www.mayoclinicproceedings.org>).

SEVERE HEADACHE AND MIGRAINE

Most of the studies we reviewed defined cervicogenic headache, migraine, and tension-type headache with criteria from the International Classification of Headache Disorders, second edition.^{109,110}

Acupuncture

Coeytaux et al¹¹¹ randomized patients with chronic daily headache to management by a neurologist with or without acupuncture. The acupuncture group had significantly reduced headache impact test scores vs the control group, but there were no significant differences in pain severity.

Massage

One small study that compared myofascial trigger point massage, a sham device, and wait list found that massage reduced headache frequency but did not significantly reduce the intensity or duration of headaches.¹¹²

Omega-3 Fatty Acids

One study randomized adolescents with migraine to omega-3 fatty acid or placebo in a crossover study.¹¹³ Adolescents experienced a reduction in headache frequency when taking either fish oil or placebo, but there was no significant difference between the treatments. Another study randomized patients with chronic headaches to increased omega-3 and lower omega-6 fatty acids in the diet or to lower omega-6 fatty acids in the diet.¹¹⁴ The participants on high omega-3 and low omega-6 fatty acid diets had greater improvement on the Headache Impact Test and in the number of headache days.

Relaxation Techniques

Six trials that we reviewed evaluated stress management, relaxation, or biofeedback for headache disorders.¹¹⁵⁻¹²⁰ Slavin-Spenney et al¹²⁰

TABLE 3. Summary of Evidence for Selected Complementary Health Approaches by Type of Pain (Sham or Placebo and/or Attention Controls)^{a,b}

Approach	Back pain	Fibromyalgia	OA of knee	Neck pain	Severe headache/migraine
Acupuncture	1 Positive trial, 2 negative	1 Positive trial, 3 negative trials	1 Positive trial, 3 negative	NA	NA
Chondroitin	NA	NA	1 Negative trial	NA	NA
Glucosamine	NA	NA	2 Positive trials, 3 negative trials	NA	NA
Chondroitin and glucosamine	NA	NA	1 Positive trial, 2 negative trials	NA	NA
Massage therapy	1 Positive trial	NA	NA	2 Positive trials	1 Positive trial
MSM	NA	NA	1 Positive trial	NA	NA
Omega-3 fatty acids	NA	NA	NA	NA	1 Negative trial
Relaxation approaches	NA	1 Positive trial, 2 negative	NA	NA	3 Positive trials
SAMe	NA	NA	NA	NA	NA
Spinal manipulation	6 Positive trials, 3 negative	NA	NA	1 Negative trial	1 Positive trial
Osteopathic manipulation	1 Positive trial, 1 negative	NA	NA	NA	NA
Tai chi	NA	2 Positive trials	3 Positive trials	NA	NA
Yoga	1 Positive trial	NA	1 Positive trial	NA	NA

^aMSM = methylsulfonylmethane; OA = osteoarthritis; NA = no US randomized controlled trials identified; SAMe = S-adenosylmethionine.

^bPositive trials are those in which the complementary approach provided statistically significant improvements in pain severity or pain-related disability or function compared with the control group. Negative trials are those in which no difference was seen between groups.

randomized students with chronic headache to expression training, relaxation training, or wait list. Both active treatments produced significant decreases in headache frequency vs the wait list control but did not differ among themselves. Two trials examined complementary approaches to either tension headaches or migraine.^{116,117} D'Souza et al¹¹⁷ randomized students to relaxation training, written emotional disclosure, or neutral writing. The relaxation group had greater reductions in headache frequency and the associated headache disability compared with the other 2 groups. Devineni and Blanchard¹¹⁶ randomized participants to an Internet behavioral intervention or a wait list. Participants in the behavioral intervention had improvement in the headache index score vs those in the wait list group. A trial randomized children with migraine to hand-warming biofeedback with stress management, hand-cooling biofeedback (attention control), or wait list.¹¹⁹ The warming biofeedback group had improved headache index scores compared with the other groups. Holroyd et al¹¹⁸ randomized participants with chronic tension headache to tricyclic antidepressant, placebo, stress management, or stress management plus tricyclic

antidepressant. Both the medication and stress management groups had improvements over placebo, but the combination produced the best outcomes. Blanchard et al¹¹⁵ randomized patients with headache to biofeedback with relaxation training, biofeedback plus cognitive therapy, sham meditation, or a headache monitoring control condition. All of the treatment groups including the sham meditation group had improvements in the headache index score in comparison with the monitoring control group.

Spinal and Osteopathic Manipulation

One trial randomized patients with chronic cervicogenic headache to 2 doses of SM or light massage.¹²¹ Based on the literature, the light massage should have little, if any, specific effects and was therefore chosen as a control for time and physical contact with the patient. Haas et al¹²¹ found improvement over all time points favoring SM compared with light massage and a dose effect with the SM. Two RCTs were identified that assessed manipulation in individuals with tension headache.^{122,123} The first trial randomized patients with chronic tension-type headache to SM or amitriptyline and found no

TABLE 4. Summary of Evidence for Selected Complementary Health Approaches by Type of Pain (Wait List, Usual Care, or Routine Care Controls)^{a,b}

Approach	Back pain	Fibromyalgia	OA of knee	Neck pain	Severe headache/migraine
Acupuncture	2 Positive trials	NA	2 Positive trials	NA	1 Positive trial
Massage therapy	3 Positive trials, 1 negative	NA	2 Positive trials	3 Positive trials	NA
Natural products supplements	NA	NA	NA	NA	NA
Relaxation approaches	NA	4 Positive trials	NA	NA	4 Positive trials
Spinal manipulation	4 Positive trials, 3 negative	NA	NA	NA	NA
Osteopathic manipulation	2 Positive trials, 2 negative	NA	NA	NA	NA
Tai chi	NA	NA	1 Positive trial	NA	NA
Yoga	4 Positive trials	1 Positive trial	NA	NA	NA

^aNo US RCTs identified; OA = osteoarthritis.

^bPositive trials are those in which the complementary approach provided statistically significant improvements in pain severity or pain-related disability or function compared with the control group. Negative trials are those in which no difference was seen between groups.

differences between the groups¹²²; however, the trial did not appear to be powered to detect noninferiority. The second study was a small trial that randomized patients to osteopathic manipulation, a palpation examination, or no treatment.¹²³ The authors noted an improvement in headache severity for the SM group; however, no statistical comparisons were made between groups. Nelson et al¹²⁴ randomized patients with migraine to amitriptyline, SM, or both treatments. Their study found no significant differences between the groups; however, it did not appear that the trial was powered to detect noninferiority.

Additional information on all headache RCTs can be found in [Supplemental Table 5](#) (available online at <http://www.mayoclinicproceedings.org>).

OVERALL SUMMARY OF RCT DATA

[Tables 3](#) and [4](#) provide concise summaries of the reviewed clinical trial data for each complementary approach stratified by painful health conditions and various control groups. In these tables, positive trials are those in which the complementary approach provided statistically significant improvements in pain severity or pain-related disability or function compared with the control group. Negative trials are those in which no difference was seen between groups. Based on a preponderance of positive trials vs negative trials, current evidence suggests that the following complementary approaches may help some patients manage their painful health conditions: acupuncture and yoga for back pain;

acupuncture and tai chi for OA of the knee; massage therapy for neck pain with adequate doses and for short-term benefit; and relaxation techniques for severe headaches and migraine. Weaker evidence suggests that massage therapy, SM, and osteopathic manipulation might also be of some benefit to those with back pain, and relaxation approaches and tai chi might help those with fibromyalgia.

SAFETY

Generally, the reporting of safety data in the reviewed RCTs was minimal. For those trials that did report safety data, we have summarized this information in the text for each painful health condition. In no case did an RCT identify a serious adverse event associated with any of the complementary approaches examined. The most common adverse events (gastrointestinal distress) were noted in trials of dietary supplements (glucosamine, chondroitin, MSM, SAME). In some trials, tai chi and yoga were associated with minor muscle or joint soreness, and acupuncture was associated with minor pain and/or bruising at the needling site.

Comparisons to Recent Systematic Reviews

Our search criteria identified a number of recent (2010 or later) systematic reviews that covered our topics of interest.¹²⁵⁻¹³³ Conclusions from these systematic reviews for practitioner approaches (acupuncture, chiropractic, massage therapy)¹²⁵⁻¹³¹ and dietary supplements¹³² were generally consistent with our findings. For instance, in a comprehensive review of both pharmacological and nonpharmacological

approaches to management of back pain, Chou et al¹²⁵ found that acupuncture and yoga appear to be effective for improving pain and/or function in patients with back pain. A Cochrane systematic review concluded that acupuncture was a viable treatment option for OA of the knee.¹²⁷ Also concurring with the present analysis, the meta-analysis by Deare et al¹²⁶ concluded that acupuncture was not an effective therapy for pain or function in individuals with fibromyalgia. In their systematic review, Posadzki and Ernst¹²⁸ found little data supporting the use of SM for headaches. Supporting our conclusions is a recent comprehensive meta-analysis of trials studying glucosamine for OA,¹³² which found considerable variability in results across trials and concluded that neither glucosamine sulfate nor glucosamine HCl provides pain relief. The RCTs examined in 2 systematic reviews of yoga for arthritis^{129,130} overlapped considerably with RCTs in the present review. The authors of the systematic reviews concluded that yoga appears to be a viable option for relieving pain and discomfort associated with arthritis but that larger, better designed trials were needed. A recent systematic review¹³¹ that included international trials found that “clinically relevant effects of OMT were found for reducing pain and improving functional status” for those with back pain. Although this conclusion is stronger than ours, the authors identified deficiencies in trial methodology and called for larger, better quality RCTs to provide firm conclusions. Not all recent systematic reviews agreed with our conclusions. A recent Cochrane Collaboration meta-analysis¹³³ concluded that SM was no more effective than “inert” interventions for managing back pain and related disability. However, this review only included RCTs published through 2009. We reviewed 8 RCTs published since then. Of these 8 later RCTs, 2 were negative trials^{27,32} and 6 were positive trials.^{36,41-43,46,54} Inclusion of these trials into the meta-analysis might have led Rubinstein et al¹³³ to draw a different conclusion.

Caveats

A number of methodological issues temper our conclusions. The trial samples tend to be white, female, and older, with very few, if any, minority group participants; as such, the generalizability of the findings to the breadth of

patients seen by primary care physicians in the United States is still unresolved. Often, the trials reviewed were small, with fewer than 100 total participants. Small trials are prone to more variability and to false-negative results. In many of the trials in which the statistical superiority of a given complementary health approach was reported, it was not clear if the differences vs the control group were clinically relevant. For the given painful health condition, a wide number of outcome measures were often used to assess pain and function. This plethora of outcomes may partly explain the conflicting results seen across trials. For most complementary approaches, there are no standard treatment protocols or algorithms, and in the case of dietary supplements, no rigorously established dosages and products; as such, trials of a given complementary approach rarely compare the exact same intervention. Our findings that relatively few mild adverse events and no serious adverse events were associated with complementary approaches are consistent with the findings from a number of systematic reviews.¹²⁵⁻¹³³ However, even large clinical trials are not powered to identify infrequent adverse events, and therefore, it is likely that this review underestimates the entire range of events associated with the complementary approaches examined. Finally, our review was intended to be an overview of data from RCTs performed in the United States. The inclusion of RCTs performed outside the United States may have resulted in a different set of recommendations.

SUPPLEMENTAL ONLINE MATERIAL

Supplemental material can be found online at <http://www.mayoclinicproceedings.org>. Supplemental material attached to journal articles has not been edited, and the authors take responsibility for the accuracy of all data.

Abbreviations and Acronyms: **ACR** = American College of Rheumatology; **cLBP** = chronic low back pain; **FIQ** = Fibromyalgia Impact Questionnaire; **HCl** = hydrochloride; **LBP** = low back pain; **MSM** = methylsulfonylmethane; **NDI** = Neck Disability Index; **OA** = osteoarthritis; **ODI** = Oswestry Disability Index; **OMT** = osteopathic manipulative therapy; **RCT** = randomized, controlled clinical trial; **SAMe** = S-adenosylmethionine; **SM** = spinal manipulation; **VAS** = visual analog scale; **WOMAC** = Western Ontario and McMaster Universities Osteoarthritis Index

Correspondence: Address to Richard L. Nahin, PhD, MPH, National Center for Complementary and Integrative Health,

National Institutes of Health, 6707 Democracy Blvd, Ste 401, Bethesda, MD 20892-5475 (NahinR@mail.nih.gov). Individual reprints of this article and a bound reprint of the entire Symposium on Pain Medicine will be available for purchase from our website www.mayoclinicproceedings.org.

The Symposium on Pain Medicine will continue in an upcoming issue.

REFERENCES

1. Nahin RL. Estimates of pain prevalence and severity in adults: United States, 2012. *J Pain*. 2015;16(8):769-780.
2. Institute of Medicine Committee on Advancing Pain Research and Education. *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research*. Washington, DC: National Academies Press; 2011.
3. Lethbridge-Cejku M, Schiller JS, Bernadel L. Summary health statistics for U.S. adults: National Health Interview Survey, 2002. *Vital Health Stat 10*. 2004;(222):1-151.
4. Pleis JR, Lucas JW. Summary health statistics for U.S. adults: National Health Interview Survey, 2007. *Vital Health Stat 10*. 2009;(240):1-159.
5. Blackwell DL, Lucas JW, Clarke TC. Summary health statistics for U.S. adults: National Health Interview Survey, 2012. *Vital Health Stat 10*. 2014;(260):1-161.
6. Walitt B, Nahin RL, Katz RS, Bergman MJ, Wolfe F. The prevalence and characteristics of fibromyalgia in the 2012 National Health Interview Survey. *PLoS One*. 2015;10(9):e0138024.
7. Barnes PM, Bloom B, Nahin RL. Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*. 2008;(12):1-23.
8. Barnes PM, Powell-Griener E, McFann K, Nahin RL. Complementary and alternative medicine use among adults: United States, 2002. *Adv Data*. 2004;(343):1-19.
9. Clarke TC, Black LI, Stussman BJ, Barnes PM, Nahin RL. Trends in the use of complementary health approaches among adults: United States, 2002-2012. *Natl Health Stat Report*. 2015;(79):1-16.
10. Eisenberg DM, Davis RB, Ettner SL, et al. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA*. 1998;280(18):1569-1575.
11. Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States: prevalence, costs, and patterns of use. *N Engl J Med*. 1993;328(4):246-252.
12. Stussman BJ, Black LI, Barnes PM, Clarke TC, Nahin RL. Wellness-related use of common complementary health approaches among adults: United States, 2012. *Natl Health Stat Report*. 2015;(85):1-12.
13. Upchurch DM, Rainisch BW. The importance of wellness among users of complementary and alternative medicine: findings from the 2007 National Health Interview Survey. *BMC Complement Altern Med*. 2015;15:362.
14. Nahin RL, Stussman BJ, Herman PM. Out-of-pocket expenditures on complementary health approaches associated with painful health conditions in a nationally representative adult sample. *J Pain*. 2015;16(11):1147-1162.
15. Cherkin DC, Eisenberg D, Sherman KJ, et al. Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med*. 2001;161(8):1081-1088.
16. Cherkin DC, Sherman KJ, Avins AL, et al. A randomized trial comparing acupuncture, simulated acupuncture, and usual care for chronic low back pain. *Arch Intern Med*. 2009;169(9):858-866.
17. Wang SM, Dezinno P, Lin EC, et al. Auricular acupuncture as a treatment for pregnant women who have low back and pos-

- terior pelvic pain: a pilot study. *Am J Obstet Gynecol*. 2009;201(3):271.e1-271.e9.
18. Wasan AD, Kong J, Pham LD, Kaptchuk TJ, Edwards R, Gollub RL. The impact of placebo, psychopathology, and expectations on the response to acupuncture needling in patients with chronic low back pain. *J Pain*. 2010;11(6):555-563.
19. Cherkin DC, Sherman KJ, Kahn J, et al. A comparison of the effects of 2 types of massage and usual care on chronic low back pain: a randomized, controlled trial. *Ann Intern Med*. 2011;155(1):1-9.
20. Eisenberg DM, Post DE, Davis RB, et al. Addition of choice of complementary therapies to usual care for acute low back pain: a randomized controlled trial. *Spine (Phila Pa 1976)*. 2007;32(2):151-158.
21. Field T, Figueiredo B, Hernandez-Reif M, Diego M, Deeds O, Ascencio A. Massage therapy reduces pain in pregnant women, alleviates prenatal depression in both parents and improves their relationships. *J Bodyw Mov Ther*. 2008;12(2):146-150.
22. Field T, Hernandez-Reif M, Diego M, Fraser M. Lower back pain and sleep disturbance are reduced following massage therapy. *J Bodyw Mov Ther*. 2007;11(2):141-145.
23. Hernandez-Reif M, Field T, Krasnegor J, Theakston H. Lower back pain is reduced and range of motion increased after massage therapy. *Int J Neurosci*. 2001;106(3-4):131-145.
24. Jacobson EE, Meleger AL, Bonato P, et al. Structural integration as an adjunct to outpatient rehabilitation for chronic nonspecific low back pain: a randomized pilot clinical trial. *Evid Based Complement Alternat Med*. 2015;2015:813418.
25. Wilson E, Payton O, Donegan-Shoaf L, Dec K. Muscle energy technique in patients with acute low back pain: a pilot clinical trial. *J Orthop Sports Phys Ther*. 2003;33(9):502-512.
26. Andersson GB, Lucente T, Davis AM, Kappler RE, Lipton JA, Leurgans S. A comparison of osteopathic spinal manipulation with standard care for patients with low back pain. *N Engl J Med*. 1999;341(19):1426-1431.
27. Cruser dA, Maurer D, Hensel K, Brown SK, White K, Stoll ST. A randomized, controlled trial of osteopathic manipulative treatment for acute low back pain in active duty military personnel. *J Man Manip Ther*. 2012;20(1):5-15.
28. Hensel KL, Buchanan S, Brown SK, Rodriguez M, Cruser dA. Pregnancy research on osteopathic manipulation optimizing treatment effects: the PROMOTE study. *Am J Obstet Gynecol*. 2015;212(1):108.e1-108.e9.
29. Licciardone JC, Buchanan S, Hensel KL, King HH, Fulda KG, Stoll ST. Osteopathic manipulative treatment of back pain and related symptoms during pregnancy: a randomized controlled trial. *Am J Obstet Gynecol*. 2010;202(1):43.e1-43.e8.
30. Licciardone JC, Minotti DE, Gatchel RJ, Keams CM, Singh KP. Osteopathic manual treatment and ultrasound therapy for chronic low back pain: a randomized controlled trial. *Ann Fam Med*. 2013;11(2):122-129.
31. Licciardone JC, Stoll ST, Fulda KG, et al. Osteopathic manipulative treatment for chronic low back pain: a randomized controlled trial. *Spine (Phila Pa 1976)*. 2003;28(13):1355-1362.
32. Bialosky JE, George SZ, Hom ME, Price DD, Staud R, Robinson ME. Spinal manipulative therapy-specific changes in pain sensitivity in individuals with low back pain (NCT01168999). *J Pain*. 2014;15(2):136-148.
33. Brennan GP, Fritz JM, Hunter SJ, Thackeray A, Delitto A, Erhard RE. Identifying subgroups of patients with acute/sub-acute "nonspecific" low back pain: results of a randomized clinical trial. *Spine (Phila Pa 1976)*. 2006;31(6):623-631.
34. Bronfort G, Evans RL, Maiers M, Anderson AV. Spinal manipulation, epidural injections, and self-care for sciatica: a pilot study for a randomized clinical trial. *J Manipulative Physiol Ther*. 2004;27(8):503-508.

35. Bronfort G, Goldsmith CH, Nelson CF, Boline PD, Anderson AV. Trunk exercise combined with spinal manipulative or NSAID therapy for chronic low back pain: a randomized, observer-blinded clinical trial. *J Manipulative Physiol Ther.* 1996;19(9):570-582.
36. Bronfort G, Hondras MA, Schulz CA, Evans RL, Long CR, Grimm R. Spinal manipulation and home exercise with advice for subacute and chronic back-related leg pain: a trial with adaptive allocation. *Ann Intern Med.* 2014;161(6):381-391.
37. Bronfort G, Maiers MJ, Evans RL, et al. Supervised exercise, spinal manipulation, and home exercise for chronic low back pain: a randomized clinical trial. *Spine J.* 2011;11(7):585-598.
38. Cherkin DC, Deyo RA, Battié M, Street J, Barlow W. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med.* 1998;339(15):1021-1029.
39. Cleland JA, Fritz JM, Kulig K, et al. Comparison of the effectiveness of three manual physical therapy techniques in a subgroup of patients with low back pain who satisfy a clinical prediction rule: a randomized clinical trial. *Spine (Phila Pa 1976).* 2009;34(25):2720-2729.
40. Cook C, Learman K, Showalter C, Kabbaz V, O'Halloran B. Early use of thrust manipulation versus non-thrust manipulation: a randomized clinical trial. *Man Ther.* 2013;18(3):191-198.
41. Fritz JM, Magel JS, McFadden M, et al. Early physical therapy vs usual care in patients with recent-onset low back pain: a randomized clinical trial. *JAMA.* 2015;314(14):1459-1467.
42. George JW, Skaggs CD, Thompson PA, Nelson DM, Gavard JA, Gross GA. A randomized controlled trial comparing a multimodal intervention and standard obstetrics care for low back and pelvic pain in pregnancy [published correction appears in *Am J Obstet Gynecol.* 2014;210(6):574-575]. *Am J Obstet Gynecol.* 2013;208(4):295.e1-295.e7.
43. Goertz CM, Long CR, Hondras MA, et al. Adding chiropractic manipulative therapy to standard medical care for patients with acute low back pain: results of a pragmatic randomized comparative effectiveness study. *Spine (Phila Pa 1976).* 2013;38(8):627-634.
44. Gudavalli MR, Cambron JA, McGregor M, et al. A randomized clinical trial and subgroup analysis to compare flexion-distraction with active exercise for chronic low back pain. *Eur Spine J.* 2006;15(7):1070-1082.
45. Haas M, Group E, Kraemer DF. Dose-response for chiropractic care of chronic low back pain. *Spine J.* 2004;4(5):574-583.
46. Haas M, Vavrek D, Peterson D, Polissar N, Neradilek MB. Dose-response and efficacy of spinal manipulation for care of chronic low back pain: a randomized controlled trial. *Spine J.* 2014;14(7):1106-1116.
47. Hadler NM, Curtis P, Gillings DB, Stinnett S. A benefit of spinal manipulation as adjunctive therapy for acute low-back pain: a stratified controlled trial. *Spine (Phila Pa 1976).* 1987;12(7):702-706.
48. Hoehler FK, Tobis JS, Buerger AA. Spinal manipulation for low back pain. *JAMA.* 1981;245(18):1835-1838.
49. Hoiris KT, Pflieger B, McDuffie FC, et al. A randomized clinical trial comparing chiropractic adjustments to muscle relaxants for subacute low back pain. *J Manipulative Physiol Ther.* 2004;27(6):388-398.
50. Hondras MA, Long CR, Cao Y, Rowell RM, Meeker WC. A randomized controlled trial comparing 2 types of spinal manipulation and minimal conservative medical care for adults 55 years and older with subacute or chronic low back pain. *J Manipulative Physiol Ther.* 2009;32(5):330-343.
51. Hsieh CY, Adams AH, Tobis J, et al. Effectiveness of four conservative treatments for subacute low back pain: a randomized clinical trial. *Spine (Phila Pa 1976).* 2002;27(11):1142-1148.
52. Hurwitz EL, Morgenstern H, Harber P, et al. A randomized trial of medical care with and without physical therapy and chiropractic care with and without physical modalities for patients with low back pain: 6-month follow-up outcomes from the UCLA low back pain study. *Spine (Phila Pa 1976).* 2002;27(20):2193-2204.
53. Pope MH, Phillips RB, Haugh LD, Hsieh CY, MacDonald L, Haldeman S. A prospective randomized three-week trial of spinal manipulation, transcutaneous muscle stimulation, massage and corset in the treatment of subacute low back pain. *Spine (Phila Pa 1976).* 1994;19(22):2571-2577.
54. Schneider M, Haas M, Glick R, Stevans J, Landsittel D. Comparison of spinal manipulation methods and usual medical care for acute and subacute low back pain: a randomized clinical trial. *Spine (Phila Pa 1976).* 2015;40(4):209-217.
55. Sutlive TG, Mabry LM, Easterling EJ, et al. Comparison of short-term response to two spinal manipulation techniques for patients with low back pain in a military beneficiary population. *Mil Med.* 2009;174(7):750-756.
56. Nyiendo J, Haas M, Goldberg B, Sexton G. Patient characteristics and physicians' practice activities for patients with chronic low back pain: a practice-based study of primary care and chiropractic physicians. *J Manipulative Physiol Ther.* 2001;24(2):92-100.
57. Saper RB, Boah AR, Keosaian J, Cerrada C, Weinberg J, Sherman KJ. Comparing once- versus twice-weekly yoga classes for chronic low back pain in predominantly low income minorities: a randomized dosing trial. *Evid Based Complement Alternat Med.* 2013;2013:658030.
58. Saper RB, Sherman KJ, Cullum-Dugan D, Davis RB, Phillips RS, Culpepper L. Yoga for chronic low back pain in a predominantly minority population: a pilot randomized controlled trial. *Altern Ther Health Med.* 2009;15(6):18-27.
59. Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann Intern Med.* 2005;143(12):849-856.
60. Sherman KJ, Cherkin DC, Wellman RD, et al. A randomized trial comparing yoga, stretching, and a self-care book for chronic low back pain. *Arch Intern Med.* 2011;171(22):2019-2026.
61. Williams K, Abildso C, Steinberg L, et al. Evaluation of the effectiveness and efficacy of iyengar yoga therapy on chronic low back pain. *Spine (Phila Pa 1976).* 2009;34(19):2066-2076.
62. Williams KA, Petronis J, Smith D, et al. Effect of iyengar yoga therapy for chronic low back pain. *Pain.* 2005;115(1-2):107-117.
63. Wolfe F, Smythe HA, Yunus MB, et al. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia: report of the Multicenter Criteria Committee. *Arthritis Rheum.* 1990;33(2):160-172.
64. Buckelew SP, Conway R, Parker J, et al. Biofeedback/relaxation training and exercise interventions for fibromyalgia: a prospective trial. *Arthritis Care Res.* 1998;11(3):196-209.
65. Assefi NP, Sherman KJ, Jacobsen C, et al. A randomized clinical trial of acupuncture compared with sham acupuncture in fibromyalgia. *Ann Intern Med.* 2005;143(1):10-19.
66. Hamis RE, Tian X, Williams DA, et al. Treatment of fibromyalgia with formula acupuncture: investigation of needle placement, needle stimulation, and treatment frequency. *J Altern Complement Med.* 2005;11(4):663-671.
67. Hamis RE, Zubieta JK, Scott DJ, Napadow V, Gracely RH, Clauw DJ. Traditional Chinese acupuncture and placebo (sham) acupuncture are differentiated by their effects on mu-opioid receptors (MORs). *Neuroimage.* 2009;47(3):1077-1085.
68. Martin DP, Sletten CD, Williams BA, Berger IH. Improvement in fibromyalgia symptoms with acupuncture: results of a randomized controlled trial. *Mayo Clin Proc.* 2006;81(6):749-757.

69. Nelson DV, Bennett RM, Barkhuizen A, et al. Neurotherapy of fibromyalgia? *Pain Med.* 2010;11(6):912-919.
70. Cash E, Salmon P, Weissbecker I, et al. Mindfulness meditation alleviates fibromyalgia symptoms in women: results of a randomized clinical trial. *Ann Behav Med.* 2015;49(3):319-330.
71. Hsu MC, Schubiner H, Lumley MA, Stracks JS, Clauw DJ, Williams DA. Sustained pain reduction through affective self-awareness in fibromyalgia: a randomized controlled trial. *J Gen Intern Med.* 2010;25(10):1064-1070.
72. Astin JA, Berman BM, Bausell B, Lee WL, Hochberg M, Forsy KL. The efficacy of mindfulness meditation plus Qigong movement therapy in the treatment of fibromyalgia: a randomized controlled trial. *J Rheumatol.* 2003;30(10):2257-2262.
73. Menzies V, Lyon DE, Elswick RK Jr, McCain NL, Gray DP. Effects of guided imagery on biobehavioral factors in women with fibromyalgia. *J Behav Med.* 2014;37(1):70-80.
74. Menzies V, Taylor AG, Bourguignon C. Effects of guided imagery on outcomes of pain, functional status, and self-efficacy in persons diagnosed with fibromyalgia. *J Altern Complement Med.* 2006;12(1):23-30.
75. Liptan G, Mist S, Wright C, Arzt A, Jones KD. A pilot study of myofascial release therapy compared to Swedish massage in fibromyalgia. *J Bodyw Mov Ther.* 2013;17(3):365-370.
76. Jones KD, Sherman CA, Mist SD, Carson JW, Bennett RM, Li F. A randomized controlled trial of 8-form Tai chi improves symptoms and functional mobility in fibromyalgia patients. *Clin Rheumatol.* 2012;31(8):1205-1214.
77. Wang C, Schmid CH, Rones R, et al. A randomized trial of tai chi for fibromyalgia. *N Engl J Med.* 2010;363(8):743-754.
78. Carson JW, Carson KM, Jones KD, Bennett RM, Wright CL, Mist SD. A pilot randomized controlled trial of the Yoga of Awareness program in the management of fibromyalgia. *Pain.* 2010;151(2):530-539.
79. Cook AJ, Wellman RD, Cherkin DC, Kahn JR, Sherman KJ. Randomized clinical trial assessing whether additional massage treatments for chronic neck pain improve 12- and 26-week outcomes. *Spine J.* 2015;15(10):2206-2215.
80. Field T, Diego M, Gonzalez G, Funk CG. Neck arthritis pain is reduced and range of motion is increased by massage therapy. *Complement Ther Clin Pract.* 2014;20(4):219-223.
81. Sherman KJ, Cherkin DC, Hawkes RJ, Miglioretti DL, Deyo RA. Randomized trial of therapeutic massage for chronic neck pain. *Clin J Pain.* 2009;25(3):233-238.
82. Sherman KJ, Cook AJ, Wellman RD, et al. Five-week outcomes from a dosing trial of therapeutic massage for chronic neck pain. *Ann Fam Med.* 2014;12(2):112-120.
83. Evans R, Bronfort G, Schulz C, et al. Supervised exercise with and without spinal manipulation performs similarly and better than home exercise for chronic neck pain: a randomized controlled trial. *Spine (Phila Pa 1976).* 2012;37(11):903-914.
84. Hurlwitz EL, Morgenstern H, Harber P, Kominski GF, Yu F, Adams AH. A randomized trial of chiropractic manipulation and mobilization for patients with neck pain: clinical outcomes from the UCLA neck-pain study. *Am J Public Health.* 2002;92(10):1634-1641.
85. Maiers M, Bronfort G, Evans R, et al. Spinal manipulative therapy and exercise for seniors with chronic neck pain. *Spine J.* 2014;14(9):1879-1889.
86. Gudavalli MR, Salsbury SA, Vining RD, et al. Development of an attention-touch control for manual cervical distraction: a pilot randomized clinical trial for patients with neck pain. *Trials.* 2015;16:259.
87. Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AM, Hochberg MC. Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial. *Ann Intern Med.* 2004;141(12):901-910.
88. Berman BM, Singh BB, Lao L, et al. A randomized trial of acupuncture as an adjunctive therapy in osteoarthritis of the knee. *Rheumatology (Oxford).* 1999;38(4):346-354.
89. Chen LX, Mao JJ, Fernandes S, et al. Integrating acupuncture with exercise-based physical therapy for knee osteoarthritis: a randomized controlled trial. *J Clin Rheumatol.* 2013;19(6):308-316.
90. Suarez-Almazor ME, Looney C, Liu Y, et al. A randomized controlled trial of acupuncture for osteoarthritis of the knee: effects of patient-provider communication. *Arthritis Care Res (Hoboken).* 2010;62(9):1229-1236.
91. Clegg DO, Reda DJ, Harris CL, et al. Glucosamine, chondroitin sulfate, and the two in combination for painful knee osteoarthritis. *N Engl J Med.* 2006;354(8):795-808.
92. Frestedt JL, Walsh M, Kuskowski MA, Zenk JL. A natural mineral supplement provides relief from knee osteoarthritis symptoms: a randomized controlled pilot trial. *Nutr J.* 2008;7:9.
93. Leffler CT, Philippi AF, Leffler SG, Masure JC, Kim PD. Glucosamine, chondroitin, and manganese ascorbate for degenerative joint disease of the knee or low back: a randomized, double-blind, placebo-controlled pilot study. *Mil Med.* 1999;164(2):85-91.
94. McAlindon T, Formica M, LaValley M, Lehmer M, Kabbara K. Effectiveness of glucosamine for symptoms of knee osteoarthritis: results from an Internet-based randomized double-blind controlled trial. *Am J Med.* 2004;117(9):643-649.
95. Messier SP, Mihalko S, Loeser RF, et al. Glucosamine/chondroitin combined with exercise for the treatment of knee osteoarthritis: a preliminary study. *Osteoarthritis Cartilage.* 2007;15(11):1256-1266.
96. Rindone JP, Hiller D, Collacott E, Nordhaugen N, Arriola G. Randomized, controlled trial of glucosamine for treating osteoarthritis of the knee. *West J Med.* 2000;172(2):91-94.
97. Rubin BR, Talent JM, Kongtawelert P, Pertusi RM, Foman MD, Gracy RW. Oral polymeric N-acetyl-D-glucosamine and osteoarthritis. *J Am Osteopath Assoc.* 2001;101(6):339-344.
98. Zenk JL, Helmer TR, Kuskowski MA. The effects of milk protein concentrate on the symptoms of osteoarthritis in adults: an exploratory, randomized, double-blind, placebo-controlled trial. *Curr Therapeutic Res.* 2002;63(7):430-442.
99. Perlman AI, Ali A, Njike VY, et al. Massage therapy for osteoarthritis of the knee: a randomized dose-finding trial. *PLoS One.* 2012;7(2):e30248.
100. Perlman AI, Sabina A, Williams AL, Njike VY, Katz DL. Massage therapy for osteoarthritis of the knee: a randomized controlled trial. *Arch Intern Med.* 2006;166(22):2533-2538.
101. Altman R, Asch E, Bloch D, et al. Development of criteria for the classification and reporting of osteoarthritis: classification of osteoarthritis of the knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism Association. *Arthritis Rheum.* 1986;29(8):1039-1049.
102. Kim LS, Axelrod LJ, Howard P, Buratovich N, Waters RF. Efficacy of methylsulfonylmethane (MSM) in osteoarthritis pain of the knee: a pilot clinical trial. *Osteoarthritis Cartilage.* 2006;14(3):286-294.
103. Najm WI, Reinsch S, Hoehler F, Tobis JS, Harvey PW. S-adenosyl methionine (SAMe) versus celecoxib for the treatment of osteoarthritis symptoms: a double-blind cross-over trial [SRCTN36233495]. *BMC Musculoskelet Disord.* 2004;5:6.
104. Brismée JM, Paige RL, Chyu MC, et al. Group and home-based tai chi in elderly subjects with knee osteoarthritis: a randomized controlled trial. *Clin Rehabil.* 2007;21(2):99-111.
105. Hartman CA, Manos TM, Winter C, Hartman DM, Li B, Smith JC. Effects of Tai Chi training on function and quality of life indicators in older adults with osteoarthritis. *J Am Geriatr Soc.* 2000;48(12):1553-1559.
106. Tsai PF, Chang JY, Beck C, Kuo YF, Keefe FJ. A pilot cluster-randomized trial of a 20-week Tai Chi program in elders with cognitive impairment and osteoarthritic knee: effects on pain and other health outcomes. *J Pain Symptom Manage.* 2013;45(4):660-669.
107. Wang C, Schmid CH, Hibberd PL, et al. Tai Chi is effective in treating knee osteoarthritis: a randomized controlled trial. *Arthritis Rheum.* 2009;61(11):1545-1553.

108. Park J, McCaffrey R, Dunn D, Goodman R. Managing osteoarthritis: comparisons of chair yoga, Reiki, and education (pilot study). *Holist Nurs Pract*. 2011;25(6):316-326.
109. Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders: 2nd edition. *Cephalalgia*. 2004; 24(suppl 1):9-160.
110. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia*. 2013;33(9):629-808.
111. Coeytaux RR, Kaufman JS, Kaptchuk TJ, et al. A randomized, controlled trial of acupuncture for chronic daily headache. *Headache*. 2005;45(9):1113-1123.
112. Moraska AF, Stenerson L, Butryn N, Krutsch JP, Schmiege SJ, Mann JD. Myofascial trigger point-focused head and neck massage for recurrent tension-type headache: a randomized, placebo-controlled clinical trial. *Clin J Pain*. 2015;31(2):159-168.
113. Harel Z, Gascon G, Riggs S, Vaz R, Brown W, Exil G. Supplementation with omega-3 polyunsaturated fatty acids in the management of recurrent migraines in adolescents. *J Adolesc Health*. 2002;31(2):154-161.
114. Ramsden CE, Faurot KR, Zamora D, et al. Targeted alteration of dietary n-3 and n-6 fatty acids for the treatment of chronic headaches: a randomized trial. *Pain*. 2013;154(11):2441-2451.
115. Blanchard EB, Appelbaum KA, Nicholson NL, et al. A controlled evaluation of the addition of cognitive therapy to a home-based biofeedback and relaxation treatment of vascular headache. *Headache*. 1990;30(6):371-376.
116. Devineni T, Blanchard EB. A randomized controlled trial of an Internet-based treatment for chronic headache. *Behav Res Ther*. 2005;43(3):277-292.
117. D'Souza PJ, Lumley MA, Kraft CA, Dooley JA. Relaxation training and written emotional disclosure for tension or migraine headaches: a randomized, controlled trial. *Ann Behav Med*. 2008;36(1):21-32.
118. Holroyd KA, O'Donnell FJ, Stensland M, Lipchik GL, Cordingley GE, Carlson BW. Management of chronic tension-type headache with tricyclic antidepressant medication, stress management therapy, and their combination: a randomized controlled trial. *JAMA*. 2001;285(17):2208-2215.
119. Scharff L, Marcus DA, Masek BJ. A controlled study of minimal-contact thermal biofeedback treatment in children with migraine. *J Pediatr Psychol*. 2002;27(2):109-119.
120. Slavin-Spenny O, Lumley MA, Thakur ER, Nevedal DC, Hijazi AM. Effects of anger awareness and expression training versus relaxation training on headaches: a randomized trial. *Ann Behav Med*. 2013;46(2):181-192.
121. Haas M, Spegman A, Peterson D, Aickin M, Vavrek D. Dose response and efficacy of spinal manipulation for chronic cervicogenic headache: a pilot randomized controlled trial. *Spine J*. 2010;10(2):117-128.
122. Bolin PD, Kassak K, Bronfort G, Nelson C, Anderson AV. Spinal manipulation vs. amitriptyline for the treatment of chronic tension-type headaches: a randomized clinical trial. *J Manipulative Physiol Ther*. 1995;18(3):148-154.
123. Hoyt WH, Shaffer F, Bard DA, et al. Osteopathic manipulation in the treatment of muscle-contraction headache. *J Am Osteopath Assoc*. 1979;78(5):322-325.
124. Nelson CF, Bronfort G, Evans R, Bolin P, Goldsmith C, Anderson AV. The efficacy of spinal manipulation, amitriptyline and the combination of both therapies for the prophylaxis of migraine headache. *J Manipulative Physiol Ther*. 1998;21(8): 511-519.
125. Chou R, Deyo R, Friedly J, et al. *Noninvasive Treatments for Low Back Pain*. Rockville, MD: Agency for Healthcare Research and Quality; 2016:Comparative Effectiveness Reviews; No. 169. AHRQ publication 16-EHC004-EF.
126. Deare JC, Zheng Z, Xue CC, et al. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev*. 2013;(5):CD007070.
127. Manheimer E, Cheng K, Linde K, et al. Acupuncture for peripheral joint osteoarthritis. *Cochrane Database Syst Rev*. 2010;(1): CD001977.
128. Posadzki P, Ernst E. Spinal manipulations for the treatment of migraine: a systematic review of randomized clinical trials. *Cephalalgia*. 2011;31(8):964-970.
129. Haaz S, Bartlett SJ. Yoga for arthritis: a scoping review. *Rheum Dis Clin North Am*. 2011;37(1):33-46.
130. Sharma M. Yoga as an alternative and complementary approach for arthritis: a systematic review. *J Evid Based Complementary Altern Med*. 2014;19(1):51-58.
131. Franke H, Franke JD, Fryer G. Osteopathic manipulative treatment for nonspecific low back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord*. 2014;15:286.
132. Wu D, Huang Y, Gu Y, Fan W. Efficacies of different preparations of glucosamine for the treatment of osteoarthritis: a meta-analysis of randomised, double-blind, placebo-controlled trials. *Int J Clin Pract*. 2013;67(6):585-594.
133. Rubinstein SM, Terwee CB, Assendelft WJ, de Boer MR, van Tulder MW. Spinal manipulative therapy for acute low back pain: an update of the Cochrane review. *Spine (Phila Pa 1976)*. 2013;38(3):E158-E177.